


EXPLANATORY DATA  
REDDING SHEET  
GEOLOGIC MAP OF CALIFORNIA  
OLAF P. JENKINS EDITION  
Compiled by Rudolph G. Strand, 1962  
(Second Printing, 1969)

INDEX TO GEOLOGIC MAPPING  
USED IN THE COMPILATION OF THE  
REDDING SHEET



- Albers, J. P., Geology of the French Gulch quadrangle, Shasta and Trinity Counties, California: U.S. Geol. Survey Bull. 1141-J in progress, Pl. 1: Geologic map of the French Gulch quadrangle, California, scale 1:62,500.
- Albers, J. P. and Robertson, J. F., 1961, Geology and ore deposits of the East Shasta copper-zinc district, Shasta County, California: U.S. Geol. Survey Prof. Paper 338, Pl. 1: Geologic map and sections of the East Shasta copper-zinc district, Shasta County, California, scale 1:24,000.
- Anonymous, Geologic maps of the west side of the Sacramento Valley, California, scale approx. 1:100,000, unpublished, 1953-55.
- Aune, Q. A., Reconnaissance geologic map of the Yolla Bolly quadrangle, scale 1:62,500, California Div. Mines and Geology, unpublished 1953-61.
- Bisio, R., Reconnaissance geology of the northeast portion of the Chancelulla Peak quadrangle and the adjacent portion of the Ono quadrangle, scale 1:62,500, California Dept. Water Resources, unpublished 1961.
- Colwell, W., De Lapp, J., and Gladish, E., 1956, Soil-vegetation map of the southeast one-fourth of the Iaqua Buttes quadrangle: U.S. Forest Service, Pl. 26 C-4: scale 1:31,680. (Adapted for state map use by W. Colwell.)
- Cox, D. P., Geology of the Helena quadrangle, Trinity County, California, scale 1:62,500, Stanford University, unpublished PhD thesis, 1956.
- Curtin, G., Geologic map, Ono quadrangle, scale 1:62,500, California Dept. Water Resources, unpublished 1959.
- Davis, G. A., Metamorphic and igneous geology of pre-Cretaceous rocks, Coffee Creek area, northeastern Trinity Alps, Klamath Mountains, California, scale 1:21,120, Univ. of California, Berkeley, unpublished PhD thesis, 1961.
- Dibblee, T. W., Jr., Geologic maps of the Scotia, Weott, Cape Mendocino, Pt. Delgada and Garberville quadrangles, scale 1:62,500, unpublished 1948-50. (Addition of fault in Shelter Cove area from mapping by S. J. Rice, California Div. Mines and Geology, 1961.)
- Diller, J. S., 1906, Redding Folio, California: U. S. Geol. Survey Geol. Atlas of the U.S., folio 138, 14 pp., Areal Geology Plate, scale 1:125,000.
- Evans, J. R., Geology of a portion of the Redding 7 1/2 minute quadrangle, scale 1:24,000: California Div. Mines and Geol., unpublished 1961.
- Evenson, R. E., 1959, Geology and ground-water features of the Eureka area, Humboldt County, California: U.S. Geol. Survey Water-Supply Paper 1470, 80 pp., Pl. 1: Generalized geologic map of Eureka area, California, showing hydrologic features, scale 1:62,500. (Modified in part by other mapping, see Rice, item 31.)
- Girard, L. V., Geological engineering study of a proposed tunnel through South Fork Mountain, Trinity County, California, scale 1:24,000, University of California, Berkeley, unpublished MS thesis, 1959.
- Harbaugh, J. W., Reconnaissance geological map of Shasta Lake area, California, scale 1:62,500, Stanford University, unpublished, 1956.
- Hicks, B., Geology of the Burnt Ranch Dam Site, Trinity County, California, scale 1:24,000, California Dept. Water Resources, unpublished 1959-60.
- Hinds, N. E. A., 1933, Geologic formations of the Redding-Weaverville districts, northern California: California Div. Mines Rept. 29, pp. 77-122, Pl. 3: Geologic map of the Redding and Weaverville quadrangles and the northwestern part of the Red Bluff quadrangle, scale 1"=4 miles.
- Hollister, V. F., Geology and ore deposits of the Shasta gold district, scale approx. 1:24,000, University of California, Berkeley, unpublished MA thesis, 1950.
- Irwin, W. P., Geology of the Weaverville quadrangle, scale 1:62,500, U. S. Geol. Survey, work in progress 1961.
- Irwin, W. P., 1960, Geologic reconnaissance of the Northern Coast Ranges and Klamath Mountains, with a summary of the mineral resources: California Div. Mines Bull. 179, 80 pp., Pl. 1 Geologic map of northwestern California (Reconnaissance mapping by William P. Irwin and Donald B. Tatlock of the U. S. Geol. Survey), scale 1:500,000. (Additional information for the Chancelulla Peak quadrangle supplied by J. I. Mallory of the U. S. Forest Service Pacific Southwest Forest and Range Experiment Station. Notes on faults in the Hyampom Valley and in Hay Fork basin from H. D. MacGinitie, 1937. The flora of the Weaverville beds of Trinity County, California, with descriptions of the plant-bearing beds: Carnegie Institution of Washington, Pub. no. 465 Part 3, pp. 83-151. Additions to geology in area just north of Hyampom from P. A. Lydon, California Div. Mines and Geology, unpublished, 1962.)
- Kinkel, A. R., Jr., Hall, W. E., and Albers, J. P., 1956, Geology and base-metal deposits of West Shasta copper-zinc district, Shasta County, California: U. S. Geol. Survey Prof. Paper 285, 156 pp., Pl. 1: Geologic map of the West Shasta copper-zinc district, Shasta County, California scale 1:24,000.
- Klein, I., and Lydon, P. A., Geological map of the southeast quarter of the Minersville quadrangle, scale 1:24,000, U. S. Bur. Reclamation, and California Div. Mines and Geology, unpublished 1953-61.
- Lipman, P. W., Geology of the southeastern Trinity Alps, California, scale approx. 1:31,000, Stanford University, PhD thesis in progress, 1961.
- Lorens, P. J., Shasta County Investigation—Appendix C, Geologic map of Shasta County, scale approx. 1" = 3 1/2 miles, California Dept. Water Resources, unpublished office report, 1955-56.
- Lydon, P. A., Reconnaissance geology of an area within the Bollibokka Mtn., Millville and Tuscan Buttes quadrangles, scale 1:62,500, California Div. Mines and Geology, unpublished, 1960.
- Manning, G. A., and Ogle, B. A., 1950, Geology of the Blue Lake quadrangle, California: California Div. Mines Bull. 148, Pl. 1: Geologic map of the Blue Lake quadrangle, California, scale 1:62,500.
- Murphy, M. A., 1956, Lower Cretaceous stratigraphic units of northern California: Am. Assoc. Petroleum Geologists Bull., v. 40, no. 9, pp. 2098-2119, Fig. 4: Geologic map of a portion of the Ono quadrangle, Shasta County, California, scale 1:62,500.
- Murphy, M. A., and Rodda, P. J., 1960, Mollusca of the Cretaceous Bald Hills Formation of California: Jour. of Paleontology, v. 34, no. 5, pp. 835-858, Fig. 1: Geologic map and columnar sections of the Bald Hills Formation, scale 13/16 inches = 1 mile.
- Ogle, B. A., 1953, Geology of Eel River Valley area, Humboldt County, California: California Div. of Mines Bull. 164, 128 pp., Pl. 1: Geologic map of the Eel River Valley area, California, scale 1:62,500.
- Olmsted, F. H., and Davis, G. H., 1961, Geologic features and ground-water storage capacity of the Sacramento Valley, California: U.S. Geol. Survey Water-Supply Paper 1497, Pl. 2: Geologic map of the Sacramento Valley, California, scale 1:250,000. (Additional information regarding faults in this area provided by J. L. Burnett, California Div. Mines and Geology, 1961. Upper Cretaceous strata in this area were mapped by H. H. Sullwold, Jr. of George H. Roth & Associates, 1960.)
- Rice, S. J., Geology of the Eureka quadrangle, California, scale 1:62,500, California Div. Mines and Geology, unpublished, 1951-52.
- Southern Pacific Company, Land Dept., Regional geologic mapping program. Geologic maps of T34-36N R1-8W and T35-36N R9W MDB & M (parts of Bollibokka Mountain, Lamoine, Schell Mountain, and Minersville quadrangles) by R. J. Ancril, H. F. Bonham, Jr., A. Chew, J. T. Collier, J. W. Cooksley, W. L. Coonrad, R. R. Dorsey, E. Kojan, R. T. Laird, W. A. Oesterling, G. W. Olcott, D. E. Pruss, R. C. Resler, M. Schafer, F. A. Stejer, J. S. Vanderpool, and R. E. Willson, scale 1:24,000, unpublished 1957-1961.
- Stinson, M. C., 1957, Geology of the Island Mountain copper mine, Trinity County, California: California Jour. Mines and Geology, v. 53, nos. 1 & 2, pp. 9-33, Pl. 1: Geologic map of the Island Mountain area, Trinity County, California, scale 1" = 1500'.
- Trask, P. D. and others, Geologic map of the Hoaglin quadrangle, scale 1:125,000, U. S. Geol. Survey, unpublished, ca. 1945-50. Modified by W. P. Irwin and D. Tatlock of the U. S. Geol. Survey 1953-55 (see item 20).
- U. S. Forest Service, Pacific Southwest Forest and Range Experiment Station State Co-operative Soil-Vegetation Survey conducted in association with the Univ. of California, for the California Division of Forestry, 1957, 1958, 1959, by K. Gowans, O. Olson, J. De Lapp, and J. I. Mallory. (Soil-vegetation maps of the Tehama County portions of the Anderson, Colyear Springs, Ono, Red Bank, and Tuscan Buttes quadrangles are available from the U. S. Forest Service as Pls. 32A 1-4, 32B 3-4, 31A 3-4, 32C 1-3, and 31D 1-4 scale 1:31,680). (Soil units adapted for state map use, and modified by field reconnaissance mapping by P. A. Lydon, California Div. Mines and Geology, 1961.)
- Voloshin, V., Reconnaissance engineering geology of the westside feeder system in Shasta and Tehama Counties, Geologic map of a portion of the Colyear Springs quadrangle, scale 1" = 1/2 mile, California Dept. Water Resources, unpublished, 1960.

For a complete list of published geologic maps of this area see Division of Mines Special Report 52.

# STRATIGRAPHIC NOMENCLATURE— REDDING SHEET

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit, are listed in stratigraphic sequence from youngest to oldest.)</small>
CENOZOIC  QUATERNARY           Pleistocene           Pliocene	Recent	<b>RECENT DUNE SAND</b>	Modern and older beach sand along the coast.
	Qs		
	Qal	<b>RECENT ALLUVIUM</b>	Alluvium, old alluvium, and young stream terrace deposits.
	Qsc	<b>RECENT RIVER AND MAJOR STREAM CHANNEL DEPOSITS IN THE GREAT VALLEY</b>	Recent sand, gravel, silt, and minor amounts of clay deposited along channels, flood plains, and natural levees of major streams.
	Qf	<b>RECENT ALLUVIAL FAN DEPOSITS IN THE GREAT VALLEY</b>	Pleistocene and Recent fan deposits which overlie the unnamed Pleistocene gravel deposits shown here as Qc, (fan deposits include Victor Formation as mapped by Olmsted and Davis, 1958).
	Qrv <sup>b</sup>	<b>RECENT VOLCANIC ROCKS:</b>  <b>BASALTIC</b>	Recent basalt flows.
	Qrv <sup>P</sup>	<b>PYROCLASTIC</b>	Recent pyroclastic deposits.
	Ql	<b>QUATERNARY LAKE DEPOSITS</b>	Pleistocene glacial lake deposits.
	Qg	<b>QUATERNARY GLACIAL DEPOSITS</b>	Pleistocene glacial moraines of the following principal glacial episodes: Late (Morris Meadow), Middle (Rush Creek), Early (Alpine Lake) and Ancient (Swift Creek). (The three youngest are judged Wisconsin, the oldest pre-Wisconsin.)
	Qt	<b>QUATERNARY NONMARINE TERRACE DEPOSITS</b>	Rohnerville Formation— <i>gravel, sand, and clay</i> . Undifferentiated terrace deposits (may be equivalent to Rohnerville or Hookton Formations).
	Qm	<b>PLEISTOCENE MARINE DEPOSITS AND MARINE TERRACE DEPOSITS</b>	Hookton Formation in the Eureka 15 minute quadrangle (includes some nonmarine units; grades laterally southward into the predominantly nonmarine Hookton shown as Qc).
	Qc	<b>PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS</b>	Hookton Formation— <i>yellow-orange gravel, sand, and clay</i> (present in Humboldt County; includes some marine shales which are dominant to north where Hookton is shown as Qm). Red Bluff Formation— <i>well-rounded boulders and gravel within a tan to brick-red iron-stained matrix of sand and some clay</i> ; unnamed gravels of basalt and andesite which unconformably overlie the Tuscan Formation and which may be stratigraphically lower than the Red Bluff Formation. Undifferentiated later terrace deposits associated with the Red Bluff Formation (Red Bank quadrangle).
	Qpv <sup>b</sup>	<b>PLEISTOCENE VOLCANIC ROCKS:</b>  <b>BASALTIC</b>	Pleistocene basalt flows.
	QP	<b>PLIOCENE-PLEISTOCENE NONMARINE SEDIMENTARY DEPOSITS</b>	Carlotta Formation— <i>massive conglomerate with sandstone and claystone</i> (marine in part).
	*	<b>QUATERNARY AND/OR PLIOCENE CINDER CONES</b>	Cinder cones of Recent age.
Puc	<b>UPPER PLIOCENE NONMARINE SEDIMENTARY ROCKS</b>	Tehama Formation— <i>poorly sorted pale yellow to greenish gray silt, silty clay, locally tuffaceous sand and gravel which weathers to pale buff and yellow-brown color</i> . Contains the Nomlaki tuff member near its base.	
Pu	<b>UPPER PLIOCENE MARINE SEDIMENTARY ROCKS</b>	Scotia Bluffs Sandstone— <i>massive sandstone containing mudstone members</i> ; Rio Dell Formation— <i>massive mudstone, alternating thin sandstone and mudstone, phantom-banded mudstone, very fine-grained sandstone, and siltstone</i> .	
Pml	<b>MIDDLE AND/OR LOWER PLIOCENE MARINE SEDIMENTARY ROCKS</b>	Eel River Formation— <i>dark gray-black mudstone, siltstone and glauconitic sandstone</i> .	
Pva	<b>PLIOCENE VOLCANIC ROCKS:</b>  <b>ANDESITIC</b>	Light to dark gray flows of medium to coarse grained hornblende or pyroxene andesite, weathers to yield subrounded boulders of fresh rock embedded in deeply weathered reddish soil.	
Pvb	<b>BASALTIC</b>	Dark gray augite-olivine basalt, medium gray porphyritic hypersthene basalt, and diabase. Basalt in the Tuscan Buttes quadrangle (Table Mtn.) which overlies the Tuscan Formation and underlies the Red Bluff Formation.	
PvP	<b>PYROCLASTIC</b>	Nomlaki Tuff Member of the Tuscan and Tehama Formations— <i>pale gray or salmon-pink, massive, poorly consolidated andesite and dacite tuff</i> , Tuscan Formation— <i>andesitic tuff, tuff breccia, agglomerate, and basal sand and gravel beds, includes dacite tuff member that is locally welded and characterized by abundant black glass</i> (much of the formation is considered to be a volcanic mudflow deposit).	

# STRATIGRAPHIC NOMENCLATURE — Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California.</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit, are listed in stratigraphic sequence from youngest to oldest.)</small>		
CENOZOIC	TERTIARY	Miocene	Mu	UPPER MIOCENE MARINE SEDIMENTARY ROCKS  Pullen Formation— <i>mudstone, in part diatomaceous, siltstone, and local basal sandstone member.</i>	
			Oc	OLIGOCENE NONMARINE SEDIMENTARY ROCKS  Weaverville Formation— <i>fine grained sandstone, shaly sandstone, sandy shale, lake beds, lignitic shale, lignite, tuff, and coarse stream conglomerate (yields reddish soil).</i>	
		Ec	EOCENE NONMARINE SEDIMENTARY ROCKS  Montgomery Creek Formation— <i>gray-green arkosic sandstone, conglomerate, and shale, light colored quartz sandstone, and thin coal seams.</i>		
		Tm	TERTIARY MARINE SEDIMENTARY ROCKS  Wildcat Group, undifferentiated— <i>mudstone, siltstone and sandstone (upper Miocene to upper Pliocene). Falor Formation—poorly cemented gray-to buff sandstone, gray to tan clay, lenticular limestone, thin red beds, and pebbly conglomerate (Pliocene).</i>		
		Ti	TERTIARY INTRUSIVE (HYPABYSSAL) ROCKS: UNDIFFERENTIATED  Porphyritic anorthoclase trachyte, shallow plug-like intrusive rock.		
	CRETACEOUS	Undivided	K	UNDIVIDED CRETACEOUS MARINE SEDIMENTARY ROCKS  Undifferentiated Cretaceous graywacke and shale of the coastal area (includes the area shown by Ogle, 1953, as the False Cape shear zone).	
			Ku	UPPER CRETACEOUS MARINE SEDIMENTARY ROCKS  Yager Formation— <i>dark gray mudstone, shale, graywacke, and conglomerate.</i> Bald Hills Formation— <i>conglomerate, sandstone, and mudstone.</i> Upper Cretaceous strata in the northern part of the Sacramento Valley— <i>gray, tan and brown well-bedded shale, sandy shale, sandstone, and some conglomerate beds near base.</i>	
			Kl	LOWER CRETACEOUS MARINE SEDIMENTARY ROCKS  Ono Formation— <i>thick mudstone, siltstone, conglomerate, graywacke, and limestone;</i> Rector Formation— <i>thin sandstone and conglomerate.</i> Lower Cretaceous strata on west side of Sacramento Valley— <i>massive conglomerate, dark-green shale, conglomeratic sandstone, and sandstone.</i> (in part this may include some Knoxville or Upper Jurassic strata in the area north of Elder Creek). Lower Cretaceous strata in the Weaverville quadrangle.	
			KJf	FRANCISCAN FORMATION  Franciscan Formation— <i>massive graywacke and minor amounts of platy, dark-gray shale, thin-bedded chert, greenstone where undifferentiated, and glaucophane schist;</i> also some areas of mildly metamorphosed equivalents of the sedimentary and volcanic rocks of the Franciscan Formation.	
			KJfv	FRANCISCAN VOLCANIC AND METAVOLCANIC ROCKS  Diabase, basalt, and agglomerate. (Mafic volcanic rocks associated with the undifferentiated Cretaceous strata of the coastal area are shown as KJfv?).	
gr			MESOZOIC GRANITIC ROCKS  Mule Mountain area intrusive rocks— <i>trondhjemite, albite granite, and some quartz diorite</i> (Late Jurassic). Shasta Bally Mountain area— <i>biotite-hornblende-quartz diorite, granodiorite (locally ranges to gabbro).</i> Also includes "birdseye" diorite and dacite porphyry dikes (Late Jurassic or Early Cretaceous). Trinity Alps area— <i>quartz diorite.</i> Helena quadrangle and Ironside Mtn. area— <i>hornblende diorite.</i> McCloud River area— <i>mafic quartz diorite and metadiorite.</i>		
bi			MESOZOIC BASIC INTRUSIVE ROCKS  Gabbro, hornblende gabbro, and banded gabbro schist. (The rocks within the Helena quadrangle shown by this unit are possibly metavolcanic.)		
ub			MESOZOIC ULTRABASIC INTRUSIVE ROCKS  Peridotite, minor pyroxenite and dunite, largely serpentized. Area 5 miles north-northwest of Trinity Dam shown as "ub" is composed of serpentine breccia, believed to be early Tertiary.		
JURASSIC			Undivided	Jk	KNOXVILLE FORMATION  Knoxville Formation— <i>shale, containing many sandstone and calcareous layers.</i>
				Ju	UPPER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS  Unnamed Upper Jurassic strata— <i>slaty and phyllitic sandstone, shale, and minor conglomerate.</i>
	Jml	MIDDLE AND/OR LOWER JURASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS  Potem Formation— <i>brown to green shaly sandstone, thin-bedded gray to green shale and gray impure limestone</i> (includes Bagley Andesite where undifferentiated). Arvison Formation— <i>chiefly marine pyroclastic rocks, buff and maroon to lavender tuffs, some shale, sandstone and andesitic flow rocks.</i>			
	JRv	JURASSIC AND/OR TRIASSIC METAVOLCANIC ROCKS  Bagley Andesite— <i>andesitic tuff, generally fine and stratified, locally agglomeratic, contains traces of marine fossils</i> (Jurassic); Bully Hill Rhyolite— <i>porphyritic and non-porphyritic silicic volcanic rocks</i> (Triassic). Quartz porphyry hypabyssal intrusive rock (possibly genetically related to the Bully Hill Rhyolite).			
	R	TRIASSIC MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS  Modin Formation— <i>andesitic tuffaceous beds containing a basal volcanic conglomerate, and gray, thin-bedded sandstone and slaty shale;</i> Brock Shale— <i>black, medium to thick-bedded shale, calcareous in part, and red toward top;</i> Hosselkus Limestone— <i>thick to thin-bedded light gray limestone;</i> Pit Formation— <i>shale, mudstone, siltstone, limestone, and lenses of tuff and tuff breccia.</i> Limestone in the Dubakella Mtn. quadrangle.			
	m ls	PRE-CRETACEOUS METAMORPHIC ROCKS UNDIFFERENTIATED, ls = LIMESTONE AND/OR DOLOMITE  Undivided Triassic and Paleozoic rocks— <i>phyllite, meta-chert and metavolcanic rocks.</i> Trinity Alps area— <i>quartz-mica schist, graphitic phyllite, and meta-chert</i> (rocks contain relict textures and possibly correlate with less metamorphosed rocks shown as "m" in the Helena quadrangle). Undivided Triassic and Paleozoic crystalline limestone shown as "ls".			
	ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS  Kerr Ranch Schist and other rock units in the northwestern part of Redding map sheet which may be more highly metamorphosed equivalents of the unnamed Upper Jurassic strata— <i>thinly foliated dark-gray quartz-sericite schist, and epidote-bearing greenschist.</i> Phyllite metachert, metagraywacke and breccia in the North Yolla Bolly Mountain area. In the southern part of the Yolla Bolly quadrangle "m" includes rocks which may be metamorphosed Franciscan rocks or may be part of an older complex— <i>phyllite, mica-quartz schist, and slate.</i>			
	mv	PRE-CRETACEOUS METAVOLCANIC ROCKS  Diabase in the Helena quadrangle.			
	UNDIVIDED	Undivided		m	PRE-CRETACEOUS METAMORPHIC ROCKS UNDIFFERENTIATED, ls = LIMESTONE AND/OR DOLOMITE  Undivided Triassic and Paleozoic rocks— <i>phyllite, meta-chert and metavolcanic rocks.</i> Trinity Alps area— <i>quartz-mica schist, graphitic phyllite, and meta-chert</i> (rocks contain relict textures and possibly correlate with less metamorphosed rocks shown as "m" in the Helena quadrangle). Undivided Triassic and Paleozoic crystalline limestone shown as "ls".
				ms	PRE-CRETACEOUS METASEDIMENTARY ROCKS  Kerr Ranch Schist and other rock units in the northwestern part of Redding map sheet which may be more highly metamorphosed equivalents of the unnamed Upper Jurassic strata— <i>thinly foliated dark-gray quartz-sericite schist, and epidote-bearing greenschist.</i> Phyllite metachert, metagraywacke and breccia in the North Yolla Bolly Mountain area. In the southern part of the Yolla Bolly quadrangle "m" includes rocks which may be metamorphosed Franciscan rocks or may be part of an older complex— <i>phyllite, mica-quartz schist, and slate.</i>

# STRATIGRAPHIC NOMENCLATURE—Continued

AGE	STATE MAP SYMBOL	STATE MAP UNIT <small>State Map Units listed here are not necessarily in stratigraphic sequence; the sequence used has been standardized for all sheets of the Geologic Map of California</small>	STRATIGRAPHIC UNITS AND CHARACTERISTIC LITHOLOGIES <small>(The formally named formations grouped within an individual State Map Unit, are listed in stratigraphic sequence from youngest to oldest.)</small>
PERMIAN	R	<b>PERMIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	McCloud Limestone— <i>light gray, very thick-bedded limestone and dolomite, locally some chert.</i>
	Rv	<b>PERMIAN METAVOLCANIC ROCKS</b>	Dekkas Andesite— <i>keratophyre, spilitic, a thin-bedded medium gray mudstone member, fossiliferous impure limestone lenses, and silicic volcanic rocks similar to Bully Hill Rhyolite (in part Triassic); Nosoni Formation—tuffaceous black mudstone and tuff, a thin basal conglomerate, and some keratophytic tuff breccia.</i>
MISSISSIPPIAN	C	<b>UNDIVIDED CARBONIFEROUS MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	Baird Formation— <i>fossiliferous mudstone, undifferentiated tuff, and small limestone lenses (deposits probably confined to Mississippian time).</i>
	Cv	<b>CARBONIFEROUS METAVOLCANIC ROCKS</b>	Porphyritic greenstone unit of the Baird Formation; Bass Mountain Diabase— <i>altered mafic lava flows and volcanic breccia (Mississippian).</i>
	CM	<b>MISSISSIPPIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	Bragdon Formation— <i>dark greenish-gray to black thinly bedded meta-shale, interstratified metamorphosed siltstone, sandstone, grit, and conglomerate in upper part, and metamorphosed local thin-bedded chert, rhyolitic tuff, and mafic volcanic rocks in the lower part. (In the French Gulch and Lamoine 15 minute quadrangles the Bragdon Formation is in thrust contact with the underlying rock units—John Albers, personal communication, 1961).</i>
DEVONIAN	D	<b>DEVONIAN MARINE SEDIMENTARY AND METASEDIMENTARY ROCKS</b>	Kennett Formation— <i>dark gray to black siliceous meta-shale, silicic tuff and tuffaceous shale in the lower part of the formation, and massive fossiliferous limestone in upper part.</i>
	Dv	<b>DEVONIAN METAVOLCANIC ROCKS</b>	Balaklala Rhyolite— <i>silicic flows interlayered with coarse and fine pyroclastic material and small intrusive bodies (similar to the Bully Hill Rhyolite).</i>
	Dv?	<b>DEVONIAN AND PRE-DEVONIAN? METAVOLCANIC ROCKS</b>	Copley Greenstone— <i>fine-grained keratophyre and some tuff beds in lower part of unit, amygdaloidal pillow lava, and fine to coarse pyroclastic material with diabase in upper part (includes some black siliceous metabasite in the Shasta Dam quadrangle).</i>
	pSs	<b>PRE-SILURIAN METASEDIMENTARY ROCKS</b>	Abrams Mica Schist— <i>recrystallized carbonate-rich schist, mica schist, and calc-silicate rocks. (A gneissic amphibolite which may possibly be a unit of the Salmon Hornblende Schist is shown as pSs?).</i>
UNDIVIDED	pSv	<b>PRE-SILURIAN METAVOLCANIC ROCKS</b>	Salmon Hornblende Schist— <i>amphibolite, hornblende schist, amphibolite hornfels, and amphibolite migmatite. (Includes a mixed rock unit of amphibolite-rich rocks and rocks similar to those shown on this map as "m".)</i>

### NOTE

<sup>1</sup> Not in stratigraphic sequence.

## TOPOGRAPHIC QUADRANGLES WITHIN THE REDDING SHEET AVAILABLE FROM THE U.S. GEOLOGICAL SURVEY 1962

